

ATMOSPHERIC SCIENCE

Warming boosts air pollution

Atmospheric conditions play an important role in driving severe air pollution events in Beijing, China. Now research finds that global warming will enhance weather conditions favouring such events, increasing the chances of severe winter-time haze in the future.

Renhe Zhang

In recent years, the adverse economic and health impacts of air pollution in China have made the abatement of such pollution a key scientific, public, and governmental concern. However, the number of winter-time (December–February) haze days in China has significantly increased in recent decades¹, primarily attributed to greater emissions of anthropogenic pollutants associated with rapid urbanization and industrialization. Despite the occurrence of severe haze events being strongly reliant on specific weather conditions², weather condition changes are thought to play a minor role in the increased incidence of haze days³. Now, writing in *Nature Climate Change*, Wenju Cai and colleagues⁴ report that conditions conducive to severe haze events in Beijing have increased in response to anthropogenic climate change, and will likely continue to do so throughout the twenty-first century.

Haze events describe times during which atmospheric concentrations of fine particles — that is, particulate matter with a diameter less than $2.5\ \mu\text{m}$ ($\text{PM}_{2.5}$) — are high enough to reduce visibility to less than 10 km. When concentrations exceed $150\ \mu\text{g m}^{-3}$, conditions can be considered severe, surpassing the level the World Health Organization defines as dangerous to human health. Severe haze events are epitomized by January 2013 conditions, wherein vast haze covered over 30 cities, peak $\text{PM}_{2.5}$ concentrations reached $500\ \mu\text{g m}^{-3}$ in Beijing, and economic impacts totalled US\$ 3.4 billion⁵.

The intensity, duration, and spatial coverage of such severe haze events have increased. In Beijing, for example, severe haze was present for 12, 18, and 25 days of the 2014–2015, 2015–2016 and 2016–2017 winter seasons, respectively. Given the grave economic and health impacts associated with haze events, it is necessary to understand what is driving apparent changes in their frequency.

With a population of more than 1.3 billion people, China has experienced considerable economic development and

corresponding urbanization over the past ~50 years. The drastic increase in energy consumption associated with these developments has not only supported a rapid growth in gross domestic product, but also a rise in national and imported emissions⁶. An increase in anthropogenic pollutants can undoubtedly be connected to enhanced haze events observed across China, particularly in large cities such as Beijing.

However, severe haze events are also driven by specific meteorological conditions that encourage stagnant air. For example, winter-time air pollution is closely related to variability in East Asian winter monsoon intensity⁷. The observed monsoonal weakening over the past few decades⁸ has reduced wind speeds around Beijing, subdued airflow transport for pollutants, and may therefore partially contribute to the increased incidence of extreme haze days.

Now, Cai *et al.* reveal a new perspective for understanding the increasing trend of haze events in Beijing⁴, proposing that anthropogenic climate change and the associated impacts on atmospheric circulation may contribute to weather conditions being more conducive to their occurrence. Specifically, they define a haze weather index (HWI) that describes whether meteorological conditions (vertical air temperature difference between the lower and upper atmosphere, lower tropospheric wind velocity, and mid-tropospheric zonal flow) are favourable for extreme haze. Using observations, they find that positive HWI values (that is, favourable haze conditions) have increased in frequency by 10% when comparing 1948–1981 with 1982–2015. Thus, climate change over the past several decades is thought to create weather conditions more favourable for triggering haze events in Beijing.




Figure 1 | Satellite-derived aerosol optical depth (AOD) over China averaged from 1 December 2016 to 31 January 2017. The AOD, ranging from non-dimensional units 0–1, is a measure of the atmospheric opacity affected by aerosols in clear sky; the higher the AOD, the more turbid the atmosphere. High pollution levels are observed throughout eastern China due to stable air and vast winter-time emissions.

How weather conditions conducive for winter-time haze will change in the future is an additional key concern. To address this issue, Cai *et al.* utilize the results of 15 models from the Coupled Model Intercomparison Project Phase 5 (CMIP5) under the RCP8.5 (high emission) scenario. They demonstrate that the frequency of all positive HWI events will increase in the future, and that this will be most notable for more extreme conditions; for example, events with a HWI higher than 1 (analogous to January 2013 conditions) will increase by 50% when comparing 1950–1999 with 2050–2099. Haze events, particularly extremes, are therefore found to occur more frequently in the future.

Due to the severity of air pollution, the Chinese government introduced the Air Pollution Prevention and Control Action Plan⁹ in 2013. The plan implemented

extreme measures to curb emissions, including controlling current pollutant sources, optimizing industrial structures, accelerating technology transformation, and increasing clean energy supply. Although the mean winter-time concentration of atmospheric pollutants over all of China has decreased by 20% in 2015 compared to 2014 levels, it is believed that 40% of this reduction may be attributed to meteorological conditions¹⁰. Indeed, in spite of stringent emission controls, severe haze days in Beijing have continued to increase, as clearly seen over the past three winters.

Therefore, from the viewpoint of Cai *et al.*, aside from controlling air pollution by limiting pollutant emissions, a global effort to slow down global warming is also urgently needed to decrease the risk of heavy air pollution in Beijing. □

Renhe Zhang is at the Institute of Atmospheric Sciences, Fudan University, Shanghai 200433, China.

e-mail: rhzhang@fudan.edu.cn

References

1. Zhang, X. *et al.* *Atmos. Environ.* **119**, 362–373 (2015).
2. Zhang, R., Li, Q. & Zhang, R. N. *Sci. China Earth Sci.* **57**, 26–35 (2014).
3. Yang, Y., Liao H. & Lou S. *J. Geophys. Res. Atmos.* **121**, 13050–13065 (2016).
4. Cai, W., Li, K., Liao, H., Wang, H. & Wu, L. *Nat. Clim. Change* <http://dx.doi.org/10.1038/nclimate3249> (2017).
5. Mu, Q. & Zhang, S. *China Environ. Sci.* **33**, 2087–2094 (2013).
6. Sun, J., Wang, J., Wei, Y., Li, Y. & Liu, M. *Int. J. Environ. Res. Public Health* **13**, 402 (2016).
7. Li, Q., Zhang, R. & Wang, Y. *Int. J. Climatol.* **36**, 346–354 (2016).
8. Huang, R., Chen, J., Wang, L. & Lin, Z. *Adv. Atmos. Sci.* **29**, 910–942 (2012).
9. Chinese State Council *Air Pollution Prevention and Control Action Plan* (Beijing: People's Publishing House, 2013).
10. Wang, X., Wang, K. & Su, L. *Sci. Rep.* **6**, 36404 (2016).

Published online: 20 March 2017